

## *Xanthorrhoea acanthostachya* (Xanthorrhoeaceae), a new species of the Perth Region, Western Australia

D. J. Bedford

National Herbarium of N.S.W., Royal Botanic Gardens, Mrs Macquarie's Road, Sydney, N.S.W. 2000

### Abstract

Bedford D. J. *Xanthorrhoea acanthostachya* (Xanthorrhoeaceae), a new species of the Perth Region, Western Australia. Nuytsia 5(2): 317-321 (1984). *Xanthorrhoea acanthostachya* is described and illustrated. It is distinguished from other *Xanthorrhoea* species in Western Australia by the combination of very elongated, prominent clusterbracts and subulate floral bracts and from *X. australis*, its nearest relative, by its scape length to spike length ratio and leaf colour. Very few examples of the species are known.

### *Xanthorrhoea acanthostachya* Bedford, sp. nov. (Figures 1-3)


*X. australi similis*, sed scapo plus minusve spicam aequanti et foliis viridibus vel minus glaucis differt. A speciebus Australiae Occidentalis bracteis fasciculorum conspicuis longatissimus ad maturitatem et bracteis floralibus subulatis distinguenda.

*Typus*: Chatfield Rd, South Western Highway, Western Australia, 24 Nov. 1982, D. J. Bedford 4 and T. D. Macfarlane (holo: NSW; iso: PERTH).

*Trunk* short to 1.5 m tall, crowns 1 to 2. *Leaves* (terminal) in more or less hemispherical crowns, 60-70 cm long, quadrate-rhombic in transverse section, 2-2.25 mm wide and 1.5-2 mm thick, green to slightly glaucous. *Leaf-base* swollen and rigid at the junction with the leaf. Scape 40-50 cm long, 7-16 mm diam. *Spike* usually more or less equal in length to scape, (20)40-50(90) cm long and 20-40 mm diam., prickly in appearance. *Cluster-bracts* very elongated, subulate in shape, dilated at the base, usually very prominent (occasionally slightly prominent), glabrous, rarely subglabrous. *Packing-bracts* (floral bracts) subulate in shape, often twisted or folded, subglabrous to glabrous (except for occasional large bracts, which have a line of hairs along the centre of the back and hairs at the margins below the tip). *Sepals* short, acute, with short beak at the tip, glabrous except for a tuft of hairs in the beak. *Petals* more or less erect at anthesis, sometimes beaked, with an adaxial proboscis, soft and membranous, glabrous except for short hairs in and around the tip, and hairs covering the proboscis. *Fruit* acute at the tip with a persistent long style-base-point, dark brown at maturity. *Seeds* dorsi-ventrally flattened, narrow ovate to ovate (Systematics Association (1962) figures 37-38), triangular in median transverse section, semi-matt black, 11-12 mm long by 4.5-5.5 mm wide, when fully mature. (Terminology as per Lee (1966a and b) and Systematics Association (1962)).

*Other specimens examined*. WESTERN AUSTRALIA: Harvey Dam Reserve, 13 Nov. 1981, T.D. Macfarlane 659 (PERTH); Keysbrook, Nov. 1900, W.V. Fitzgerald NSW 154569 (NSW); 5 miles E of Mogumber, 25 Aug. 1970, K.M. Allan s.n. (spirit collection only) (PERTH).

*Distribution*. At present *X. acanthostachya* is known only from four sites in the Perth Region of W.A. as defined by Marchant and Perry (1981).



NSW

10 cm

2. Base + thickness 1.7 mm  
 ◇  
 green (maybe bluish)  
 53 cm  
 87 cm  
 16 mm  
 36 mm  
 1.6  
 subulate  
 100  
 most long glumes to subulate, a few shorter  
 and have silicles of hairs along the centre of the  
 mid ribs of hairs on the sides below the  
 green

44 cm (2-headed)  
 DJB 4  
 24. xi. 1982  
 Western Australia

FLORA OF AUSTRALIA PROJECT  
*Xanthorrhoea acanthostachya*  
 Beckford  
 Det. Beckford  
 Date 15. 11. 1982

NATIONAL HERBARIUM OF NEW SOUTH WALES (NSW)  
 ROYAL BOTANIC GARDENS SYDNEY

*Xanthorrhoea acanthostachya* Beckford  
 Det. D. Beckford  
 15. 11. 1982

Loc. South Western Highway  
 nr. Chaffield Rd

T. 32° 23' S Long 115° 59' E Alt. m  
 State Western Australia Subdiv

Coll. D. Beckford 4 and  
 T. J. Meesteriana Date 24. xi. 1982

N. Occasional on flat ground  
 on grey sand with thin silty gravel  
 in flat woodland  
 coastal plain near Earling sump  
 with *Leptocarpus bromeliifolius*  
 (acaulis) and *Kingia* with woolly  
 leaf bases close to  
 Plant two-headed (just)

Figure 1. *Xanthorrhoea acanthostachya*, holotype specimen in herb. NSW.




Figure 2. *Xanthorrhoea acanthostachya*, close-up view of immature spike, showing very prominent cluster-bracts. K. M. Allan s.n., 5 miles E of Mogumber, W. Austral., 25 Aug. 1970.

*Ecology.* The species occurs on the coastal plain in grey sand overlain by lateritic gravel, often with *Dasypogon bromeliifolius* and *Kingia australis*, and on steep slopes in stony lateritic soil in Jarrah (*Eucalyptus marginata*) woodland.



Figure 3. *Xanthorrhoea acanthostachya*, mature flowering spike showing exceptionally long prominent cluster-bracts. T. D. Macfarlane 659, sheet 1.

*Flowering period.* Young flowering spikes have been collected in August and flowering and recently fruiting spikes have been collected in November.

*Affinities.* There are no obvious allies of *Xanthorrhoea acanthostachya* in Western Australia, although its leaves are at least superficially similar to those of *X. preissii*. *X. australis* of eastern Australia has similarly shaped bracts and leaf cross-sectional shape, and is therefore probably the most closely allied species. *X. semiplana* of South Australia has similarly shaped floral bracts to *X. acanthostachya* but is a much more massive plant with large broad transverse-rhombic median transverse section leaves.

*Xanthorrhoea acanthostachya* differs from *X. australis* in (a) scape length to spike length ratio; *X. acanthostachya* has scape length more or less equal to spike length, *X. australis* always has a much shorter scape than spike (less than  $\frac{1}{2}$  the length), (b) leaf colour; *X. acanthostachya* has green to slightly glaucous leaves, *X. australis* has very glaucous leaves, (c) sepals; *X. acanthostachya* has short, acute sepals, with a short beak at the tip, without a proboscis in the beak, *X. australis* has subulate shaped sepals with a long narrow beak at the tip, often with a proboscis in the beak, (d) petals; *X. acanthostachya* petals sometimes have a beak, *X. australis* petals never have a beak.

*Etymology.* The specific epithet is from the Greek *akantha*, meaning a thorn or prickle, and *stakhys*, an ear of grain or a spike, in reference to the distinctly thorny or prickly appearance of the spike due to the prominent cluster-bracts.

#### Acknowledgements

I wish to thank Dr Terry Macfarlane for help with my field studies on *Xanthorrhoea* in Western Australia, Mrs Karen Wilson for advice with the Latin diagnosis and Ms Anna-louise Quirico for technical assistance. The research and field work were undertaken with the aid of a grant from the Australian Biological Resources Study.

#### References

- Lee, A. T. (1966a). Contr. New South Wales Natl. Herb. 4(3): 35-54.
- Lee, A. T. (1966b). Contr. New South Wales Natl. Herb. Fl. Ser. 34: 1-16.
- Marchant, N. G. and Perry, G. (1981). A checklist of the vascular plants of the Perth Region, Western Australia. W. Austral. Herb. Res. Notes No. 5: 111-136.
- Systematics Association (1962). Terminology of simple symmetrical plane shapes. Taxon 11(3): 145-148.

